

Estimation of Thyroid Doses Received by the Population of Belarus as a Result of the Chernobyl Accident*

Yuri GAVRILIN, Valeri KHROUCH, and Sergei SHINKARYOV
Institute of Biophysics, 46 Zhivopisnaya St., 123182 Moscow, Russia

Vladimir DROZDOVITCH, Viktor MINENKO, and Elena SHEMYAKINA
Institute of Radiation Medicine, 23 Masherova Ave., 220600 Minsk, Belarus

André BOUVILLE
National Cancer Institute, 6130 Executive Blvd., Rockville, MD 20854, USA

Lynn ANSPAUGH
Lawrence Livermore National Laboratory, 7000 East Ave., Livermore, CA 94550, USA

Abstract. Within weeks of the Chernobyl accident, about 300,000 measurements of human thyroidal ^{131}I content were conducted in the more contaminated territories of the Republic of Belarus. Results of these and other measurements form the basis of thyroid-dose reconstruction for residents of Belarus. Preliminary estimates of thyroid doses have been divided into three classes:

Class 1 ("measured" doses). Individual doses are estimated directly from the measured thyroidal ^{131}I content of the person considered, plus information on lifestyle and dietary habits. Such estimates are available for about 130,000 individuals from the contaminated areas of Gomel and Mogilev Oblasts.

Class 2 ("passport" doses). For every settlement with a sufficient number of residents with "measured" doses, individual-thyroid-dose distributions are determined for several age groups and levels of milk consumption. This action has been called the "passportization" of the settlement. A population of about 2.5 million people resides in the "passportized" settlements.

Class 3 ("inferred" doses). For any settlement where the number of residents with "measured" doses is small or equal to zero, individual thyroid doses are derived from the relationship obtained between the mean-adult-thyroid dose and the deposition density of ^{131}I or ^{137}Cs in settlements with "passport" doses presenting characteristics similar to those of the settlement considered. This method can be applied to the remainder of the population (about 7.5 million people).

An approximate estimate of the collective thyroid dose for the residents of Belarus is presented. Illustrative results of individual-thyroid dose and associated uncertainty are discussed for rural settlements and urban areas.

* Portions of this work were performed at the Lawrence Livermore National Laboratory under the auspices of the Office of International Health Studies; Assistant Secretary for Environment, Safety and Health; US Department of Energy; under contract W-7405-Eng-48.